

Anti-Vitronectin (human)**Mouse monoclonal antibody**

Subclass: IgG2b/k

CAT. NO.

CSI 003-02

Clone: HV2

SPECIFICITY	CSI 003-02 is highly specific for vitronectin. There is no evidence for cross-reactivity with other connective tissue proteins (fibronectin, elastin, collagen, laminin).
IMMUNOGEN	Human vitronectin purified from plasma by heparin-affinity chromatography
TESTED APPLICATIONS	ELISA, WB, IHC, AP
SPECIES REACTIVITY (POSITIVE)	Human
SPECIES REACTIVITY (NEGATIVE)	Cat, dog, cow, sheep, goat, pig, rabbit, horse
EPITOPE SPECIFICITY	The epitope is located in the connecting region. Binding can be competed selectively by a peptide comprising aa 121-133.

PRESENTATION

Content:	Available in 400 µL and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.
Preparation:	Protein-A purified
Form:	Liquid
Solvent:	0.01 M phosphate buffer, pH 7.4, with 0.5 M NaCl and 15 mM sodium azide
Storage:	4-8°C without exposure to light. No precautions necessary during handling.

APPLICATION

ELISA: It binds equally well to native and denatured vitronectin and can be used to quantitate vitronectin in human plasma in a sandwich ELISA with CSI 003-08. It does not interfere with the binding of any known vitronectin ligands. CSI 003-02 also binds to vitronectin in ELISA when vitronectin is coated directly onto the microtiter well. In Western blotting a dilution guideline of 1/100 has proved successful (1, 2).

WB: CSI 003-02 can be used in Western blotting (1, 2)

IHC: CSI 003-02 can be used in IHC. Please consult www.proteinatlas.org

AP: CSI 003-02 can be used to purify vitronectin from human plasma by affinity chromatography. It can also be used to quantitatively affinity-deplete human plasma or serum of vitronectin. It partially denatures vitronectin upon antibody binding.

TARGET

Vitronectin is a plasma glycoprotein that circulates in the blood. Vitronectin is circulating as a mixture of both 75 kDa and 65 kDa forms. Vitronectin is a major cell adhesive glycoprotein and is a common component of extracellular matrix and plasma. It competes effectively with other plasma proteins and is often involved in cell attachment, regulation of blood coagulation and immune responses. It has similar tissue distribution to fibronectin and also its integrin receptor recognizes fibronectin (2).

REFERENCES

- Morris CA, Underwood PA, Bean PA, Sheehan M, Charlesworth JA (1994) Relative topography of biologically active domains of human vitronectin. Evidence from monoclonal antibody epitope and denaturation studies. *J Biol Chem* 269:23845-23852.
- Underwood PA, Kirkpatrick A, Mitchell SM (2002) New insights into heparin binding to vitronectin: studies with monoclonal antibodies. *Biochem J* 365:57-67.

CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its products.