

## Anti-Fibronectin (bovine, human, chicken)

CAT. NO. CSI 005-17

**OVERVIEW** 

Product Name Anti-Fibronectin (bovine, human, chicken) Conjugation Unconjugated

DescriptionMouse monoclonal antibodyHostMouseIsotypeIgG1/kCloneA17

Tested Applications ELISA, WB, IHC, IP

**SPECIFICITY** 

Specificity CSI 005-17 is highly specific for fibronectin. There is no evidence for cross-reactivity with other

connective tissue proteins (vitronectin, elastin, collagen, laminin). The antibody inhibits integrin-

mediated cell adhesion to the cell binding domain of fibronectin.

Epitope is located in the 120kD cell binding fragment

**Immunogen** Lysed bovine corneal endothelial cells and **Gene ID** 280794, 2335, 396133

extracellular matrix

Target Fibronectin is an adhesive glycoprotein with a molecular mass of 440 kDa. It is believed to be important for

the formation of a provisional matrix that promotes cell adhesion and migration during wound healing. Its age-dependent increase in plasma and tissues may be accompanied in pathological states, especially in tumor growth, by its proteolytic breakdown by a number of neutral proteases. It has also shown that several

of its proteolytic breakdown products exhibit unexpected and mostly harmful biological activities. (1)

Species Reactivity Bovine, Human, Chicken Species Reactivity Not determined

POSITIVE NEGATIVE

**PROPERTIES** 

Form Liquid Unit Size 0,4 mL and 1 mL

**Concentration** 1 mg/mL ±15%, See CoA for lot details

Purification Protein A or Protein G purified Purification Notes BSA free

Storage buffer 0.01 M phosphate buffer, pH 7.4, with 0.5 M NaCl and 15 mM sodium azide

**Storage condition** 2-8°C without exposure to light

Safety Wear protective clothing

**TESTED APPLICATIONS** 

**ELISA** CSI 005-17 can be used in ELISA. (1, 2, 3, 4, 5)

WB In Western blotting a dilution guideline of 1/100 has proved successful. (1)

IHC CSI 005-17 can be used in immunostaining of frozen PLP-fixed sections of bovine and human tissues.IP CSI 005-17 can be used in immunoprecipitation. It can be used to probe fibronectin conformation.

## **SCIENTIFIC REFERENCES**

- 1. Underwood PA, Dalton BA, Steele JG, Bennett FA, Strike P (1992) Anti-fibronectin antibodies that modify heparin binding and cell adhesion: evidence for a new cell binding site in the heparin binding region. J Cell Sci 102:833-845.
- 2. Underwood PA, Steele JG, Dalton BA (1993) Effects of polystyrene surface chemistry on biological activity of solid phase fibronectin and vitronectin, analysed with monoclonal antibodies. J Cell Sci 104:793-803.
- 3. Di Girolamo N, Underwood PA, McCluskey PJ, Wakefield D (1993) Functional activity of plasma fibronectin in patients with Diabetes mellitis. Diabetes 42:1606-1613.

## CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/5 be responsible for loss of profits or indirect consequential losses resulting from use of its products.



- 4. Dalton BA, McFarland CD, Underwood PA, Steele JG (1995) Role of heparin binding domain of fibronectin in attachment and spreading of human bone derived cells. J Cell Sci 108:2083-2092.
- 5. Underwood PA, Bean PA, Mitchell SM, Whitelock JM (2001) Specific affinity depletion of cell adhesion molecules and growth factors from serum. J Immunol Methods 247:217-224.